

ALLANTOIN-CONTAINING SKIN CREAM

by

Elliott Farber

5

CROSS REFERENCES

This application is a continuation-in-part application of Application Serial No. 09/570,266, entitled "Allantoin-Containing Skin Cream," filed May 12, 2000 by Elliott Farber, which in turn is a continuation-in-part application of Application Serial No. 09/360,095, entitled
10 "Oil-in-Water Emulsion With Improved Stability," filed July 23, 1999 by Elliott Farber. Both of these prior applications are hereby incorporated in their entirety by this reference.

BACKGROUND OF THE INVENTION

15

This invention is directed to a skin cream containing allantoin in an oil-in-water emulsion with improved stability.

Allantoin is a commonly used ingredient in cosmetic applications, particularly for
20 skin creams, where it exerts a skin protective function. Many such cosmetic compositions and other compositions are prepared as emulsions, particularly oil-in-water emulsions. One emulsifier system used with such compositions is a combination of sodium lauryl sulfate and beeswax. Although solutions of sodium lauryl sulfate are alkaline with an approximate pH of 9.5, the simultaneous use of beeswax with its organic acids produces a complex and neutralized
25 system with a pH of about 6.8 to about 7.5. However, in such a system with a pH range of 6.8 to 7.5, allantoin degrades significantly with time and in accelerated stability tests at 40°C. Because cosmetics and other preparations designed for application to the skin are typically stored by users at room temperature, and room temperatures can fluctuate with climatic conditions, such a degree of stability is undesirable. Therefore, there is a need for an oil-in-water emulsified
30 composition containing allantoin in which the stability of allantoin is increased.

In particular, there is a need for compositions that are suitable for treating a number of severe and difficult-to-treat skin conditions. One of these skin conditions is epidermolysis bullosa. This is a severe genetic skin disorder in which the skin breaks down and large blisters appear. These blisters are difficult to treat by conventional means. Other skin diseases for which improved treatments are needed are pressure ulcers, decubitus ulcers or bed sores, and diabetic ulcers, as well as milia. Therefore, improved compositions that are suitable for treating these diseases are needed.

SUMMARY

In general, a composition according to the present invention comprises an oil-in-water emulsion comprising:

(1) allantoin; and

(2) at least one anionic or nonionic emulsifier that is substantially hydrophilic and is soluble in water.

The pH of the composition is in a range of from about 3.0 to about 6.0. This pH range stabilizes the allantoin and makes storage of the composition for extended periods practical. Preferably, the pH of the composition is in a range of from about 5.0 to about 6.0.

Typically, the at least one anionic or nonionic emulsifier is selected from the group consisting of:

(1) an acidic anionic polymer;

(2) an anionic emulsifier selected from the group consisting of sodium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, sodium dodecylbenzenesulfonate, and sodium lauryl sarcosinate;

(3) a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose chain length ranges from 8 to 22 carbon atoms;

(4) glyceryl stearate;

(5) cetyl alcohol;

- (6) stearic acid;
- (7) sodium stearyl lactylate;
- (8) sodium isostearyl lactylate;
- (9) triethanolamine;
- 5 (10) a polyethylene glycol ether of cetaryl alcohol wherein the number of polyethylene glycol moieties in the ether is from 6 to 40; and
- (11) beeswax.

10 The composition can further comprise a carbohydrate polymer selected from the group consisting of galactoarabinan, polygalactose, and polyarabinose. Typically, the carbohydrate polymer is galactoarabinan.

One embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

- 15 (1) allantoin; and
- (2) an emulsifier system comprising:
 - (a) an acidic anionic polymer; and
 - (b) a polyethylene glycol ester of stearic acid.

20 The pH of the composition is from about 3.0 to about 6.0. Preferably, the pH of the composition is from about 5.0 to about 6.0.

Typically, in this embodiment, the acidic anionic polymer is a carboxypolymethylene polymer.

25 The composition can further comprise a carbohydrate polymer. Typically, the carbohydrate polymer is selected from the group consisting of galactoarabinan, polygalactose, and polyarabinose. Preferably, the carbohydrate polymer is galactoarabinan.

30 Another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin; and

(2) an emulsifier system comprising:

(a) an acidic anionic polymer; and

(b) an anionic emulsifier that is substantially hydrophilic and is soluble in
5 water, the pH of the composition being adjusted to a range of from about 3.0 to about 6.0.
Preferably, the pH of the composition is adjusted to a range of from about 5.0 to about 6.0.

The anionic emulsifier can be selected from the group consisting of sodium lauryl
sulfate, sodium laureth sulfate, sodium oleyl succinate, sodium dodecylbenzenesulfonate, and
10 sodium lauryl sarcosinate. Preferably, the anionic emulsifier is sodium lauryl sulfate.

The composition can further comprise a carbohydrate polymer selected from the
group consisting of galactoarabinan, polygalactose and polyarabinose. Preferably, the
carbohydrate polymer is galactoarabinan.

15 Another embodiment of the present invention is a composition comprising an oil-
in-water emulsion comprising:

(1) allantoin;

(2) an emulsifier system comprising an acidic anionic polymer; and

20 (3) a base to adjust the pH of the composition to a value in a range from about 3.0
to about 6.0. Preferably, the pH of the composition is from about 5.0 to about 5.5.

25 Preferably, the acidic anionic polymer is a carboxypolymethylene polymer.
Preferably, the base is an organic base such as triethanolamine.

In another embodiment, the composition comprises an oil-in-water emulsion
comprising:

(1) allantoin;

(2) an emulsifier system comprising:

30 (a) an acidic anionic polymer; and

(b) a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose carbon chain length ranges from 8 to 22 carbon atoms, wherein the pH of the composition is from about 3.0 to about 6.0. Preferably, the pH of the composition is from about 5.0 to about 6.0.

5

In this embodiment, the emulsifier system can further comprise glyceryl stearate.

This embodiment also can include a carbohydrate polymer as described above.

10

In yet another embodiment of the present invention, a composition comprising an oil-in-water emulsion comprising:

(1) allantoin;

(2) an emulsifier system comprising:

15

(a) sodium stearoyl lactylate;

(b) sodium isostearoyl lactylate;

(c) optionally, triethanolamine;

(d) optionally, at least one nonionic emulsifier selected from the group consisting of a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose carbon chain length ranges from 8 to 22 carbon atoms; and

20

(3) an acid to adjust the pH to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from 5.0 to about 5.8.

Typically, the acid is citric acid.

25

Still another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin; and

30

(2) an emulsifier system comprising at least one polyethyleneglycol ether of cetearyl alcohol, wherein the number of polyethylene glycol moieties in the polyethyleneglycol ether of cetearyl alcohol is from 6 to 40; and

(3) an acid to adjust the pH of the composition to a range of from about 5.0 to about 5.8.

For this embodiment, the acid is typically also citric acid.

For this embodiment, the emulsifier system typically comprises cetareth-25 and cetareth-6.

Yet another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin;

(2) an emulsifier system comprising:

(a) a polyethylene glycol ester of stearic acid; and

(b) glyceryl stearate; and

(3) an acid to adjust the pH of the composition to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8.

Typically, in this embodiment, the number of ethylene glycol moieties in the polyethylene glycol ester of stearic acid is from 25 to 100. Preferably, the polyethylene glycol ester of stearic acid is PEG-100 stearate. Typically, in this embodiment, the acid is citric acid.

Still another embodiment of the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin;

(2) a carbohydrate polymer; and

(3) an emulsifier system comprising:

(a) beeswax; and

(b) an anionic emulsifier that is substantially hydrophilic and is soluble in water;

wherein the pH of the composition is in a range from about 3.0 to about 6.0. Preferably, the pH is in the range of from about 5.0 to about 6.0.

The carbohydrate polymer in this embodiment is as described above.

Typically, the anionic emulsifier that is substantially hydrophilic and soluble in
5 water is selected from the group consisting of sodium lauryl sulfate, sodium laureth sulfate,
sodium oleyl succinate, sodium dodecylbenzenesulfonate, and sodium lauryl sarcosinate.
Preferably, the anionic emulsifier is sodium lauryl sulfate.

10 The composition can further comprise citric acid to adjust the pH.

For all of these embodiments, the composition can comprise one or more
additional ingredients as described below.

The composition can comprise an emollient component comprising at least one
15 emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod
liver oil.

The composition can comprise an emollient such as butylated hydroxytoluene.

20 The composition can comprise herbal extracts such as one or more of St. John's
wort extract, witch hazel extract, chamomile extract, and arnica extract.

The composition can comprise a preservative component comprising at least one
preservative selected from the group consisting of methylparaben, propylparaben, and
25 diazolidinyl urea.

The composition can comprise a chelating agent such as tetrasodium EDTA.

The composition can comprise a solvent component comprising at least one
30 solvent selected from the group consisting of propylene glycol, butylene glycol and glycerin.
Preferably, the solvent component is propylene glycol.

Ranges of compositions are disclosed for each of these embodiments.

5

DESCRIPTION

A composition according to the present invention comprises an oil-in-water emulsion comprising:

- (1) allantoin; and
- 10 (2) at least one anionic or nonionic emulsifier that is substantially hydrophilic and is soluble in water.

The composition can further include other ingredients, such as a chelating agent to bind metal ions that might accelerate degradation of the composition. A particularly preferred
15 chelating agent is EDTA. The EDTA can be added in various acid or salt forms depending on the pH of the composition, such as EDTA itself, disodium EDTA, or tetrasodium EDTA.

The pH of the composition is in a range of from about 3.0 to about 6.0. This pH range stabilizes the allantoin and makes storage of the composition for extended periods
20 practical. Preferably, the pH is in a range of from about 5.0 to about 6.0.

Typically, the at least one anionic or nonionic emulsifier is selected from the group consisting of:

- (1) an acidic anionic polymer;
- 25 (2) an anionic emulsifier selected from the group consisting of sodium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, sodium dodecylbenzenesulfonate, and sodium lauryl sarcosinate;
- (3) an nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose chain length ranges from 8 to 22 carbon atoms;
- 30 (4) glyceryl stearate;
- (5) cetyl alcohol;

- (6) stearic acid;
- (7) sodium stearyl lactylate;
- (8) sodium isostearyl lactylate;
- (9) triethanolamine;
- 5 (10) a polyethylene glycol ether of cetearyl alcohol wherein the number of polyethylene glycol moieties in the ether is from 6 to 40; and
- (11) beeswax.

10 The composition can further comprise a carbohydrate polymer selected from the group consisting of galactoarabinan, polygalactose, and polyarabinose. Typically, the carbohydrate polymer is galactoarabinan.

One embodiment of an improved composition containing allantoin is a composition that comprises an oil-in-water emulsion comprising:

- 15 (1) allantoin; and
- (2) an emulsifier system comprising:
 - (a) an acidic anionic polymer; and
 - (b) a polyethylene glycol ester of stearic acid.

20 The pH of the composition is adjusted to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 6.0. The pH is adjusted with sodium hydroxide or other base as required.

25 The acidic anionic polymer is preferably a carboxypolymethylene polymer. Such polymers are marketed under the brand names "Carbomer" and "Carbopol." A suitable carboxypolymethylene polymer is marketed by B.F. Goodrich under the brand name "Carbomer." This is a slightly cross-linked polyacrylic acid that is from 1% to 2% cross-linked by allylsucrose or allylpentaerythritol with the polyacrylic acid. The resulting molecular weight range of this polymer is from about 2×10^6 daltons to about 1×10^9 daltons. The average
30 molecular weight of this polymer is about 4×10^6 daltons.

Preferably, the concentration of the carboxypolymethylene polymer is from about 0.5 percent to about 2 percent of the composition.

5 The composition can further comprise a carbohydrate polymer. Preferably, the carbohydrate polymer is galactoarabinan. Galactoarabinan is derived from trees of the genus Larix (larch) and is a hemicellulosic product easily extractable by water in a pure form. Galactoarabinan has been consumed by humans in common foods such as carrots, tomatoes, maple syrup, soybeans, and wheat flour, among others. The molecular weight of the galactoarabinan is about 20,000. A suitable source of galactoarabinan is Larex, Inc (White Bear
10 Lake, MN). Typically, the composition contains from about 1 percent to about 25 percent of galactoarabinan. Preferably, the composition contains from about 2 percent to about 10 percent of the carbohydrate polymer.

This embodiment of a composition according to the present invention can further
15 include other ingredients. For example, the composition can include an emollient component for smoothness. The emollient component can include at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

The composition can also include an antioxidant to prevent rancidity of
20 ingredients such as cod liver oil. A preferred antioxidant is butylated hydroxytoluene (BHT).

The composition can further include a solvent component. Typically, the solvent component can include at least one solvent selected from the group consisting of propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.
25

The composition can further include a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben, propylparaben, and diazolidinyl urea. Preferably, the preservative component comprises methylparaben, propylparaben, and diazolidinyl urea.
30

The composition can further include fragrance. The use of fragrance is well known in the cosmetic art and in the art of over-the-counter drug formulation, and many suitable fragrances are known in the art. The stability and function of the cream is not altered by the presence or absence of fragrance.

5

Optionally, this embodiment of the composition can further include herbal extracts. The herbal extracts can include one or more of St. John's wort extract, witch hazel extract, chamomile extract, and arnica extract. However, these herbal extracts are typically omitted in this embodiment.

10

This embodiment of the composition can optionally further include other components, such as proteins, humectants, other preservatives, essential oils, other vitamins, colorants, hydroxyacids, other plant extracts, chelators, sunscreens, sodium hyaluronate, lipids, fatty acids, thickeners, panthenol, and the like. The use of such components is conventional in the cosmetic art and in the over-the-counter drug art. Typical sunscreens are octyl methoxycinnamate and benzophenone-3.

15

The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention with a pH range of from about 5.0 to about 6.0.

20

Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 85.0% of this embodiment of the composition. An optimum concentration of water in this embodiment of the composition is about 69.95%.

25

The carboxypolymethylene polymer can comprise from about 0.30% to about 3.0% of this embodiment of the composition. Preferably, the carboxypolymethylene polymer comprises from about 0.50% to about 2.0% of this embodiment of the composition. An optimum concentration of the carboxypolymethylene polymer is about 0.85% of this embodiment of the composition.

30

Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

PEG-100 stearate can comprise from about 0.25% to about 2.5% of this embodiment of the composition. Preferably, PEG-100 stearate comprises from about 0.50% to about 2.0% of this embodiment of the composition. An optimum concentration of PEG-100 stearate is about 1.50% of this embodiment of the composition.

Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

Cetyl alcohol can comprise from about 1.0% to about 8.0% of this embodiment of the composition. A preferred concentration of cetyl alcohol is from about 2.0% to about 7.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 4.20% of this embodiment of the composition.

Stearyl alcohol can comprise from about 0.5% to about 6.0% of this embodiment of the composition. A preferred concentration of stearyl alcohol is from about 0.75% to about 5.0% of this embodiment of the composition. An optimum concentration of stearyl alcohol is about 1.50% of this embodiment of the composition.

Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.20% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

5

Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. A preferred concentration of methylparaben is from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

10

Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

15

Diazolidinyl urea can comprise from about 0.05% to about 0.25% of this embodiment of the composition. Preferably, diazolidinyl urea comprises from about 0.10% to about 0.20% of this embodiment of the composition. An optimum concentration of diazolidinyl urea is about 0.15% of this embodiment of the composition.

20

Allantoin can comprise from about 0.50% to about 2.0% of this embodiment of the composition. A preferred concentration of allantoin is from about 1.0% to about 2.0% of this embodiment of the composition. An optimum concentration of allantoin is about 1.50% of this embodiment of the composition.

25

Fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of the composition. As indicated above, fragrance can be omitted, and it may be desirable to omit fragrance in circumstances in which the composition is intended for use on sensitive individuals or individuals who may undergo an allergic reaction to fragrance.

30

Triethanolamine can comprise from about 0.05% to about 3.0% of this embodiment of the composition to adjust the pH. A preferred concentration of triethanolamine is from about 0.20% to about 2.0% of this embodiment of the composition. An optimum
5 concentration of triethanolamine is about 0.80% of this embodiment of the composition.

In another alternative embodiment of the composition, the emulsifier can be an anionic emulsifier that is substantially hydrophilic and is soluble in water. In this embodiment, the anionic emulsifier replaces the polyethylene glycol ester of stearic acid. This embodiment
10 further includes the acidic anionic polymer such as carboxypolymethylene. Optionally, but preferably, this alternative embodiment of the composition includes the carbohydrate polymer such as galactoarabinan.

The anionic emulsifier that is substantially hydrophilic and soluble in water can
15 be selected from the group consisting of sodium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, sodium dodecylbenzenesulfonate, and sodium lauryl sarcosinate. A particularly preferred anionic emulsifier is sodium lauryl sulfate.

Commercially available preparations of sodium lauryl sulfate contain sufficient
20 excess sodium hydroxide so that they have a pH of about 10.0. This sodium hydroxide can be used to adjust the pH when the anionic emulsifier is sodium lauryl sulfate; in this alternative, no additional alkali may be needed. When another anionic emulsifier is used, additional alkali may be required to adjust the pH.

25 In yet another alternative embodiment of the composition, the emulsifier system comprises the acidic anionic polymer as described above and a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose carbon chain length ranges from 8 to 22 carbon atoms.

30 Preferably, the acidic anionic polymer is carboxypolymethylene as described above.

This alternative embodiment of the composition can further include glyceryl stearate in the emulsifier system.

5 This embodiment of the composition has a pH from about 3.0 to 6.0, adjusted as necessary, typically with an acid. The acid can be an organic acid, an inorganic acid, or a mixture of both. Preferably, the pH is from about 5.0 to about 6.0.

10 This embodiment of the composition can further comprise a carbohydrate polymer such as galactoarabinan as described above.

15 In this embodiment of the composition, preferred organic acids include organic acids whose carbon chain length ranges from 2 to 22 carbon atoms and can be monocarboxylic, dicarboxylic, or tricarboxylic acids. The acids can be aliphatic or aromatic. Particularly preferred organic acids include citric acid, ascorbic acid, glycolic acid, lactic acid, benzoic acid, and salicylic acid. A most particularly preferred organic acid is citric acid.

20 Typically, in this embodiment of the composition, the inorganic acid is a strong acid. It can be a monoprotic, diprotic, or triprotic acid. Particularly preferred inorganic acids include hydrochloric acid, sulfuric acid, and phosphoric acid.

25 This alternative embodiment of the composition can further include other ingredients as described above, including an emollient component, an antioxidant, a solvent component, a chelating agent, herbal extracts, a preservative, and fragrance.

30 The composition can further include other components, such as proteins, humectants, other preservatives, essential oils, other vitamins, colorants, hydroxyacids, other plant extracts, sunscreens, sodium hyaluronate, lipids, fatty acids, thickeners, panthenol, and the like. The use of such components is conventional in the cosmetic art and in the over-the-counter drug art. Typical sunscreens are octyl methoxycinnamate and benzophenone-3.

In yet another embodiment of the composition, the emulsifier system comprises the acidic anionic polymer described above; one example of this acidic anionic polymer is marketed as Carbomer. In this embodiment, the pH is adjusted with an organic or inorganic base to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.5. A preferred organic base is triethanolamine. A preferred inorganic base is sodium hydroxide. In general, it is preferred to use an organic base such as triethanolamine.

This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one solvent selected from the group consisting of lanolin oil, cetyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, and cod liver oil.

This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene.

This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben and propylparaben. Preferably, the preservative component comprises both methylparaben and propylparaben.

This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention where the pH is from about 5.0 to about 5.5.

5 Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 80.0% of this embodiment of the composition. An optimum concentration of water is about 73.55% of this embodiment of the composition.

10 The carboxypolymethylene polymer can comprise from about 0.40% to about 3.0% of this embodiment of the composition. Preferably, the carboxypolymethylene polymer comprises from about 0.5% to about 2.0% of this embodiment of the composition. An optimum concentration of the carboxypolymethylene polymer is about 1.00% of the composition.

15 Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, the propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of the propylene glycol is about 5.70% of this embodiment of the composition.

20 Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.0% of this embodiment of the composition.

25 Cetyl alcohol can comprise from about 1.0% to about 8.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.0% to about 7.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 3.00% of this embodiment of the composition.

30 Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this

embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

5 Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.30% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

10 Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

15 Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

20 Allantoin can comprise from about 0.50% to about 2.0% of this embodiment of the composition. Preferably, allantoin comprises from about 1.0% to about 2.0% of this embodiment of the composition. An optimum concentration of allantoin is about 1.50% of this embodiment of the composition.

25 Fragrance, if present, can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, if present, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance, if present, is about 0.20% of this embodiment of the composition.

30 Triethanolamine, as a 95% solution, can comprise from about 0.05% to about 3.0% of this embodiment of the composition to adjust the pH to a value in the range of from about 5.0 to about 5.5. Preferably, triethanolamine comprises from about 0.20% to about 2.0%

of this embodiment of the composition to adjust the pH as indicated. An optimum concentration of triethanolamine is about 0.80% of the composition to adjust the pH as indicated.

Yet another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

- (1) allantoin; and
- (2) an emulsifier system comprising:
 - (a) cetyl alcohol; and
 - (b) stearic acid.

In this embodiment, the pH is adjusted to a range of from about 3.0 to about 6.0 by addition of a quantity of a weak organic base. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8. The weak organic base can be an amine-containing base such as ethanolamine, diethanolamine, or triethanolamine. A preferred organic base is triethanolamine.

This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one solvent selected from the group consisting of lanolin oil, cetyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, and cod liver oil.

This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene.

This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group

consisting of methylparaben and propylparaben. Preferably, the preservative component comprises both methylparaben and propylparaben.

5 This embodiment of the composition can further include fragrance as described above. The stability and function of the cream is not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention
10 when the pH is in the range of from about 5.0 to about 5.8.

Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 85.0% of this embodiment of the composition. An optimum concentration of water is about 71.70% of this embodiment of
15 the composition.

Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene
20 glycol is about 5.70% of this embodiment of the composition.

Triethanolamine can comprise from about 0.2% to about 4.0% of this embodiment of the composition. Preferably, triethanolamine comprises from about 0.5% to about 3.0% of this embodiment of the composition. An optimum concentration of
25 triethanolamine is about 1.25% of this embodiment of the composition.

Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of
30 this embodiment of the composition.

Cetyl alcohol can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.0% to about 6.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 3.50% of this embodiment of the composition.

5

Stearic acid can comprise from about 0.50% to about 5.0% of this embodiment of the composition. Preferably, stearic acid comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of stearic acid is about 2.50% of this embodiment of the composition.

10

Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.50% to about 5.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.0% of this embodiment of the composition.

15

Butylated hydroxytoluene can comprise from about 0.1% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.2% to about 0.8% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

20

Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

25

Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

30

Allantoin can comprise from about 0.50% to about 2.0% of this embodiment of the composition. Preferably, allantoin comprises from about 1.0% to about 2.0% of this embodiment of the composition. An optimum concentration of allantoin is about 1.50% of this embodiment of the composition.

5

If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

10

Still another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin; and

(2) an emulsifier system comprising:

15

(a) sodium stearoyl lactylate;

(b) sodium isostearoyl lactylate;

(c) optionally, triethanolamine;

(d) optionally, at least one nonionic emulsifier selected from the group consisting of a nonionic emulsifier that is an ethoxylated ether or an ethoxylated ester whose carbon chain length ranges from 8 to 22 carbon atoms.

20

Sodium stearoyl lactylate is the sodium salt of the stearic acid ester of lactyl lactate. Sodium isostearoyl lactylate is the sodium salt of the isostearic acid ester of lactyl lactate.

25

In this embodiment of the composition, the composition further comprises an acid to adjust the pH to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8. The acid can be an inorganic or an organic acid as described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

30

This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

5

This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, and cod liver oil.

10

This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben and propylparaben. Preferably, the preservative component comprises both methylparaben and propylparaben.

15

This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene.

20

This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

25

The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is in the range of from about 5.0 to about 5.8.

30

Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 80.0% of this embodiment

of the composition. An optimum concentration of water is about 73.72% of this embodiment of the composition.

5 Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

10 Citric acid can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, citric acid comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of citric acid is about 0.18% of this embodiment of the composition.

15 Sodium stearoyl lactylate can comprise from about 0.30% to about 3.0% of this embodiment of the composition. Preferably, sodium stearoyl lactylate comprises from about 0.50% to about 2.50% of this embodiment of the composition. An optimum concentration of sodium stearoyl lactylate is about 1.00% of this embodiment of the composition.

20 Sodium isostearoyl lactylate can comprise from about 0.05% to about 1.0% of this embodiment of the composition. Preferably, sodium isostearoyl lactylate comprises from about 0.10% to about 0.70% of this embodiment of the composition. An optimum concentration of sodium isostearoyl lactylate is about 0.25% of this embodiment of the composition.

25 Tetrasodium EDTA can comprise from about 0.05% to about 0.25% of this embodiment of the composition. Preferably, tetrasodium EDTA comprises from about 0.10% to about 0.20% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

30 Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this

embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

5 Cetyl alcohol can comprise from about 1.0% to about 8.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.0% to about 7.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 3.80% of this embodiment of the composition.

10 Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.0% of this embodiment of the composition.

15 Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.20% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

20 Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

25 Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

30 Allantoin can comprise from about 0.50% to about 2.0% of this embodiment of the composition. Preferably, allantoin comprises from about 1.0% to about 2.0% of this

embodiment of the composition. An optimum concentration of allantoin is about 1.50% of this embodiment of the composition.

If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

Still another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin; and

(2) an emulsifier system comprising at least one polyethyleneglycol ether of cetearyl alcohol.

In polyethylene glycol ethers of cetearyl alcohol suitable for use in compositions according to the present invention, the number of ethylene glycol moieties can range from 6 to 40, e.g., $R(\text{OCH}_2\text{CH}_2)_{25}\text{OH}$ where $R = \text{CH}_3(\text{CH}_2)_{16-18}$. In one preferred embodiment of the present invention, the emulsifier system comprises both cetareth-25 and cetareth-6, i.e., polyethylene glycol ethers of cetearyl alcohol with 25 and 6 ethylene glycol units respectively.

In this embodiment of the composition, the composition further comprises an acid to adjust the pH to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8. The acid can be an inorganic or an organic acid as described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

This embodiment of the invention can further comprise an emollient component.

5 The emollient component can comprise at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

10 This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene.

This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben, propylparaben, and diazolidinyl urea. Preferably, the preservative
15 component comprises all of methylparaben, propylparaben, and diazolidinyl urea.

This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.
20

The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is adjusted to a range of from about 5.0 to about 5.8.

25 Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 55.0% to about 75.0% of this embodiment of the composition. An optimum concentration of water is about 66.33% of this embodiment of the composition.

30 Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.2% to

about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

5 Tetrasodium EDTA can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, tetrasodium EDTA comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

10 Cetareth-25 can comprise from about 0.50% to about 4.0% of this embodiment of the composition. Preferably, cetareth-25 comprises from about 2.0% to about 3.5% of this embodiment of the composition. An optimum concentration of cetareth-25 is about 2.60% of this embodiment of the composition.

15 Citric acid can comprise from about 0.04% to about 0.40% of this embodiment of the composition. Preferably, citric acid comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of citric acid is about 0.12% of this embodiment of the composition.

20 Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

25 Cetyl alcohol can comprise from about 3.0% to about 10.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 3.5% to about 7.5% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 4.30% of this embodiment of the composition.

30 Stearyl alcohol can comprise from about 1.0% to about 5.0% of this embodiment of the composition. Preferably, stearyl alcohol comprises from about 2.0% to about 4.0% of this

embodiment of the composition. An optimum concentration of stearyl alcohol is about 3.50% of this embodiment of the composition.

5 Ceteareth-6 can comprise from about 0.5% to about 4.0% of this embodiment of the composition. Preferably, ceteareth-6 comprises from about 1.0% to about 3.0% of this embodiment of the composition. An optimum concentration of ceteareth-6 is about 1.80% of this embodiment of the composition.

10 Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

15 Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.20% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

20 Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

25 Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

30 Diazolidinyl urea can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, diazolidinyl urea comprises from about 0.10% to

about 0.30% of this embodiment of the composition. An optimum concentration of diazolidinyl urea is about 0.15% of this embodiment of the composition.

5 Allantoin can comprise from about 0.50% to about 2.0% of this embodiment of the composition. Preferably, allantoin comprises from about 1.0% to about 2.0% of this embodiment of the composition. An optimum concentration of allantoin is about 1.50% of this embodiment of the composition.

10 If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

15 Yet another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin; and

(2) an emulsifier system comprising:

(a) a polyethylene glycol ester of stearic acid; and

(b) glyceryl stearate.

20

Typically, the number of ethylene glycol moieties in the polyethylene glycol ester of stearic acid is from 25 to 100. Two preferred polyethylene glycol esters of stearic acid for use in this embodiment of compositions according to the present invention are PEG-40 stearate and PEG-100 stearate, with 40 and 100 ethylene glycol moieties respectively. A particularly
25 preferred polyethylene glycol ester of stearic acid is PEG-100 stearate.

In this embodiment of the composition, the composition further comprises an acid to adjust the pH to a range of from about 3.0 to about 6.0. Preferably, the pH is adjusted to a range of from about 5.0 to about 5.8. The acid can be an inorganic or an organic acid as
30 described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component is one or more of propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one emollient selected from the group consisting of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene.

This embodiment of the invention can further comprise a preservative component. The preservative component can comprise at least one preservative selected from the group consisting of methylparaben, propylparaben, and diazolidinyl urea. Preferably, the preservative component comprises all of methylparaben, propylparaben, and diazolidinyl urea.

This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is in the range of from about 5.0 to about 5.8.

Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 55.0% to about 80.0% of this embodiment of the composition. An optimum concentration of water is about 67.86% of this embodiment of the composition.

5

Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.30% to about 7.00% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

10

Tetrasodium EDTA can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, tetrasodium EDTA comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

15

Citric acid can comprise from about 0.04% to about 0.40% of this embodiment of the composition. Preferably, citric acid comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of citric acid is about 0.14% of this embodiment of the composition.

20

PEG-100 stearate can comprise from about 1.0% to about 5.0% of this embodiment of the composition. Preferably, PEG-100 stearate comprises from about 1.50% to about 3.00% of this embodiment of the composition. An optimum concentration of PEG-100 stearate is about 2.60% of this embodiment of the composition.

25

Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

30

Cetyl alcohol can comprise from about 2.0% to about 10.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.50% to about 7.50% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 3.00% of this embodiment of the composition.

5

Stearyl alcohol can comprise from about 1.0% to about 4.0% of this embodiment of the composition. Preferably, stearyl alcohol comprises from about 1.0% to about 3.5% of this embodiment of the composition. An optimum concentration of stearyl alcohol is about 2.50% of this embodiment of the composition.

10

Glyceryl stearate can comprise from about 1.0% to about 5.0% of this embodiment of the composition. Preferably, glyceryl stearate comprises from about 2.0% to about 4.0% of this embodiment of the composition. An optimum concentration of glyceryl stearate is about 2.50% of this embodiment of the composition.

15

Cod liver oil can comprise from about 1.0% to about 7.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

20

Butylated hydroxytoluene can comprise from about 0.10% to about 1.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.20% to about 0.80% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

25

Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

30

Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

5

Diazolidinyl urea can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, diazolidinyl urea comprises from about 0.10% to about 0.30% of this embodiment of the composition. An optimum concentration of diazolidinyl urea is about 0.20% of this embodiment of the composition.

10

Allantoin can comprise from about 0.50% to about 2.0% of this embodiment of the composition. Preferably, allantoin comprises from about 1.0% to about 2.0% of this embodiment of the composition. A preferred concentration of allantoin is about 1.50% of this embodiment of the composition.

15

If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, fragrance comprises from about 0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

20

Yet another embodiment of a composition according to the present invention is a composition comprising an oil-in-water emulsion comprising:

(1) allantoin;

(2) a carbohydrate polymer; and

(3) an emulsifier system comprising:

(a) beeswax; and

(b) an anionic emulsifier that is substantially hydrophilic and is soluble in

water.

25

The carbohydrate polymer is typically selected from the group consisting of galactoarabinan, polygalactose, and polyarabinose. Preferably, the carbohydrate polymer is galactoarabinan.

5 The anionic emulsifier that is substantially hydrophilic and soluble in water can be selected from the group consisting of sodium lauryl sulfate, sodium laureth sulfate, sodium oleyl succinate, sodium dodecylbenzenesulfonate, and sodium lauryl sarcosinate. A particularly preferred anionic emulsifier is sodium lauryl sulfate.

10 The pH of the composition is adjusted to a range of between about 3.0 and about 6.0, typically with an acid. Preferably, the pH is adjusted to a range of from about 5.0 to about 6.0. The acid can be an inorganic or an organic acid as described above. Preferably, the acid is a weak organic acid. Most preferably, the acid is citric acid.

15 This embodiment can further comprise other ingredients. For example, this embodiment of the invention can further comprise a solvent component. Typically, the solvent component comprises at least one solvent selected from the group consisting of propylene glycol, glycerin, or butylene glycol. Preferably, the solvent component is propylene glycol.

20 This embodiment of the invention can further comprise a chelator component. Preferably, the chelator component is tetrasodium ethylenediaminetetraacetic acid.

 This embodiment of the invention can further comprise an emollient component. The emollient component can comprise at least one emollient selected from the group consisting
25 of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil. Preferably, the emollient component comprises all of lanolin oil, cetyl alcohol, stearyl alcohol, and cod liver oil.

 This embodiment of the composition can also include an antioxidant. A preferred antioxidant is butylated hydroxytoluene.

30

This embodiment of the invention can further comprise a preservative component. The preservative component can comprise one or more of methylparaben or propylparaben. Preferably, the preservative component comprises methylparaben and propylparaben.

5 This embodiment of the composition can further include fragrance as described above. The stability and function of the cream are not altered by the presence or absence of fragrance. As indicated above, it may be desirable to omit fragrance in some cases.

10 The following discussion describes ranges, preferred concentrations and optimum concentrations for preferred compositions according to this embodiment of the present invention when the pH is adjusted to a range of from about 5.0 to about 6.0.

15 Water can comprise from about 50.0% to about 90.0% of this embodiment of the composition. Preferably, water comprises from about 60.0% to about 80.0% of this embodiment of the composition. An optimum concentration of water is about 61.65% of this embodiment of the composition.

20 Propylene glycol can comprise from about 2.0% to about 9.0% of this embodiment of the composition. Preferably, propylene glycol comprises from about 4.0% to about 7.0% of this embodiment of the composition. An optimum concentration of propylene glycol is about 5.70% of this embodiment of the composition.

25 Sodium lauryl sulfate, as a 30% solution, can comprise from about 0.50% to about 5.0% of this embodiment of the composition. Preferably, sodium lauryl sulfate, as a 30% solution, comprises from about 1.0% to about 3.0% of this embodiment of the composition. An optimum concentration of sodium lauryl sulfate, as a 30% solution, is about 1.90% of this embodiment of the composition.

30 Tetrasodium EDTA can comprise from about 0.05% to about 0.30% of this embodiment of the composition. Preferably, tetrasodium EDTA comprises from about 0.10% to

about 0.20% of this embodiment of the composition. An optimum concentration of tetrasodium EDTA is about 0.15% of this embodiment of the composition.

5 Galactoarabinan can comprise from about 1.0% to about 25.0% of this embodiment of the composition. Preferably, galactoarabinan comprises from about 3.0% to about 15.0% of this embodiment of the composition. An optimum concentration of galactoarabinan is about 5.00% of this embodiment of the composition.

10 Citric acid can comprise from about 0.05% to about 0.25% of this embodiment of the composition. Preferably, citric acid comprises from about 0.10% to about 0.20% of this embodiment of the composition. An optimum concentration of citric acid is about 0.15% of this embodiment of the composition.

15 Lanolin oil can comprise from about 5.0% to about 15.0% of this embodiment of the composition. Preferably, lanolin oil comprises from about 8.0% to about 12.0% of this embodiment of the composition. An optimum concentration of lanolin oil is about 10.60% of this embodiment of the composition.

20 Cetyl alcohol can comprise from about 1.0% to about 8.0% of this embodiment of the composition. Preferably, cetyl alcohol comprises from about 2.0% to about 7.0% of this embodiment of the composition. An optimum concentration of cetyl alcohol is about 4.20% of this embodiment of the composition.

25 Stearyl alcohol can comprise from about 0.50% to about 6.0% of this embodiment of the composition. Preferably, stearyl alcohol comprises from about 1.0% to about 4.0% of this embodiment of the composition. An optimum concentration of stearyl alcohol is about 2.00% of this embodiment of the composition.

30 Beeswax can comprise from about 0.50% to about 5.0% of this embodiment of the composition. Preferably, beeswax comprises from about 1.0% to about 3.0% of this

embodiment of the composition. An optimum concentration of beeswax is about 1.90% of this embodiment of the composition.

5 Cod liver oil can comprise from about 0.50% to about 15.0% of this embodiment of the composition. Preferably, cod liver oil comprises from about 1.0% to about 10.0% of this embodiment of the composition. An optimum concentration of cod liver oil is about 2.00% of this embodiment of the composition.

10 Butylated hydroxytoluene can comprise from about 0.1% to about 3.0% of this embodiment of the composition. Preferably, butylated hydroxytoluene comprises from about 0.25% to about 2.50% of this embodiment of the composition. An optimum concentration of butylated hydroxytoluene is about 0.50% of this embodiment of the composition.

15 Methylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, methylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of methylparaben is about 0.30% of this embodiment of the composition.

20 Propylparaben can comprise from about 0.10% to about 0.50% of this embodiment of the composition. Preferably, propylparaben comprises from about 0.15% to about 0.40% of this embodiment of the composition. An optimum concentration of propylparaben is about 0.25% of this embodiment of the composition.

25 Allantoin can comprise from about 0.50% to about 2.0% of this embodiment of the composition. Preferably, allantoin comprises from about 1.0% to about 2.0% of this embodiment of the composition. An optimum concentration of allantoin is about 1.50% of this embodiment of the composition.

30 If present, fragrance can comprise from about 0.05% to about 0.50% of this embodiment of the composition. Preferably, if present, fragrance can comprise from about

0.10% to about 0.40% of this embodiment of the composition. An optimum concentration of fragrance is about 0.20% of this embodiment of the composition.

5 Examples of particularly preferred compositions according to the present invention are described below.

10 The composition is prepared by standard mixing techniques, such as are conventional in the cosmetic art and in the art of over-the-counter drug formulation for blending lipid-soluble components and water-soluble components. These mixing techniques include both manual and mechanical mixing, and include homogenization mixing and sweep mixing. The mixing techniques to be used can be chosen by one of ordinary skill in the art based on variables such as the viscosity of the components to be mixed and the volume of those components, as well as the relative proportion of lipid-soluble and water-soluble ingredients. The composition can be mixed in two or more batches, such as one batch containing lipid-soluble ingredients and
15 another batch containing water-soluble ingredients, and the batches can then be mixed at the final stage of preparation.

The batches to be used are shown in Tables 1-11.

20 Compositions according to the present invention can be formulated for the treatment of skin diseases and conditions. Among the skin diseases and conditions for which compositions according to the present invention can be formulated are epidermolysis bullosa, decubitus ulcers, pressure ulcers, diabetic ulcers, and milia. Compositions according to the present invention can be formulated for treatment of other skin diseases and conditions. The details of the composition can be varied according to the particular condition to be treated. For
25 example, greater or lesser degrees of oil or lipid-soluble components can be included, and, in the case of compositions intended to be used on patients who may undergo allergic reactions, compounds that often generate allergic reactions, such as fragrance and coloring, can be excluded. Such details can readily be ascertained by one of ordinary skill in the art.

30

The invention is illustrated by the following Examples. These Examples are for illustrative purposes only and are not intended to limit the invention.

Example 1

5 Preparation of Skin Protectant Over-the-Counter Cream with pH of 7.4

(Prior Art Example)

A skin protectant over-the-counter (OTC) cream was prepared in accordance with the formulation of Table 1.

TABLE 1**COMPOSITION OF ALLANTOIN-CONTAINING SKIN CREAM WITH pH OF 7.4**

INGREDIENT	RANGE	PREFERRED	OPTIMUM
Part A			
Water	50.0-90.0	55.0-75.0	66.20
Sodium Lauryl Sulfate (30%)	0.50-2.50	1.00-2.50	1.90
Propylene Glycol	2.0-9.0	3.0-6.0	5.30
Tetrasodium EDTA	0.05-0.50	0.10-0.30	0.15
Part B			
Lanolin Oil	5.0-15.0	8.0-12.0	10.60
Cetyl Alcohol	3.0-10.0	3.5-7.5	6.80
Stearyl Alcohol	1.0-5.0	1.0-3.0	2.00
Beeswax	0.50-2.50	1.0-2.5	1.90
Cod Liver Oil	1.0-7.0	1.0-4.0	2.00
BHT	0.10-1.00	0.20-0.80	0.50
Part C			
St. John's Wort Extract	0.05-0.50	0.05-0.15	0.10
Witch Hazel Extract	0.05-0.50	0.05-0.15	0.10
Chamomile Extract	0.05-0.50	0.05-0.15	0.10
Arnica Extract	0.05-0.50	0.05-0.15	0.10
Methylparaben	0.10-0.50	0.15-0.40	0.30
Propylparaben	0.10-0.50	0.10-0.30	0.25
Allantoin	0.50-2.00	0.50-2.00	1.50
Fragrance	0.05-0.50	0.10-0.30	0.20

The Part A ingredients were combined and heated to 175°F with mixing. The Part B ingredients were combined and heated to 175°F with mixing. The Part B mixture was then added to the Part A mixture with mixing. The resulting mixture was then cooled to 120°F with continued mixing. The Part C ingredients were then added with mixing. The final emulsion was allowed to cool with continued mixing. The resulting cream had a pH of 7.4. Samples of the cream prepared from Example 1 were used for accelerated aging stability studies and analyzed for their allantoin concentration after a period of time at 40°C. The results are shown in Table 2.

As can be seen from Table 2, the allantoin in the cream from Example 1 undergoes degradation and would not meet the specifications required for an OTC drug.

TABLE 2**STABILITY OF ALLANTOIN IN SKIN CREAM COMPOSITION
OF EXAMPLE 1 WITH STORAGE AT 40°C**

Days at 40°C	Weight % Allantoin
0	1.5
30	1.4
60	1.3
90	1.2

Example 2Preparation of a Cream Containing Allantoin with Lower pH

An OTC skin cream containing allantoin was prepared using the ingredients in Table 3 to provide a cream with a lower pH.

TABLE 3
COMPOSITION OF ALLANTOIN-CONTAINING
SKIN CREAM WITH pH OF 5.3

INGREDIENT	RANGE	PREFERRED	OPTIMUM
Part A			
Water	50.0-90.0	55.0-75.0	68.68
Sodium Lauryl Sulfate (30%)	0.50-2.50	1.00-2.50	1.90
Propylene Glycol	2.0-9.0	3.0-6.0	5.30
Tetrasodium EDTA	0.05-0.50	0.10-0.30	0.15
Citric Acid	0.05-0.50	0.08-0.35	0.12
Part B			
Lanolin Oil	5.0-15.0	8.0-12.0	10.60
Cetyl Alcohol	3.0-10.0	3.5-7.5	4.20
Stearyl Alcohol	1.0-5.0	1.0-3.0	2.00
Beeswax	0.50-2.50	1.0-2.5	1.90
Cod Liver Oil	1.0-7.0	1.0-4.0	2.00
BHT	0.10-1.00	0.20-0.80	0.50
Part C			
St. John's Wort Extract	0.05-0.50	0.05-0.15	0.10
Witch Hazel Extract	0.05-0.50	0.05-0.15	0.10
Chamomile Extract	0.05-0.50	0.05-0.15	0.10
Arnica Extract	0.05-0.50	0.05-0.15	0.10
Methylparaben	0.10-0.50	0.15-0.40	0.30
Propylparaben	0.10-0.50	0.10-0.30	0.25
Allantoin	0.50-2.00	0.50-2.00	1.50
Fragrance	0.05-0.50	0.10-0.30	0.20

The Part A ingredients were combined and heated to 175°F with mixing. The Part B ingredients were combined and heated to 175°F with mixing. The Part B mixture was added to the Part A mixture with mixing. The resulting mixture was then cooled to 120°F with mixing at which time the Part C ingredients were added with mixing. The final emulsion was allowed to cool with continue mixing. The resulting cream had a pH of 5.3.

5

It was found that a similar cream was produced if Part B was added to Part A or Part A was added to Part B. However, the cream has a better appearance if the oil phase and water phase are homogenized under high shear after the two phases are added to one another.

Samples of the cream of this example were used for accelerated aging stability studies and analyzed for their allantoin concentration. The results are shown in Table 4. As can be seen from Table 4, the allantoin is stable over time in a cream with a pH of 5.3.

TABLE 4
STABILITY OF ALLANTOIN IN SKIN CREAM COMPOSITION OF
EXAMPLE 2 WITH STORAGE AT 40°C

Days at 40°C	Weight % Allantoin
0	1.4
30	1.4
60	1.4
90	1.4

Example 3Preparation of Allantoin-Containing Skin Cream with Ionic Emulsifiers

5 An allantoin-containing skin cream with ionic emulsifiers is prepared according to
Table 5. The preparation follows the method used in Example 2, with the ingredients in each of
Part A, Part B, and Part C being combined separately and then Part B being added to Part A, with
Part C then being added to the combination of Part A and Part B. The pH is adjusted to a value
in a range of from about 5.0 to about 5.8 by neutralizing the stearic acid with enough
10 triethanolamine to reach this pH. Other bases can be used instead of triethanolamine.

TABLE 5
ALLANTOIN-CONTAINING SKIN CREAM WITH IONIC EMULSIFIERS

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	60.0 - 85.0	71.70
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
Triethanolamine (99%)	0.20 - 4.0	0.50 - 3.0	1.25
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	1.0 - 7.0	2.0 - 6.0	3.50
Stearic Acid	0.50 - 5.0	1.0 - 4.0	2.50
Cod Liver Oil	1.0 - 7.0	1.5 - 5.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Allantoin	0.50 - 2.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20

Example 4Preparation of Allantoin-Containing Skin Cream with Lactylate Emulsifiers

5

An allantoin-containing skin cream with the emulsifiers sodium stearoyl lactylate and sodium isostearoyl lactylate is prepared according to Table 6. The preparation follows the method used in Example 3. The pH is adjusted by the addition of the appropriate quantity of citric acid.

TABLE 6
ALLANTOIN-CONTAINING SKIN CREAM WITH LACTYLATE EMULSIFIERS

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	60.0 - 80.0	73.42
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
Citric Acid	0.05 - 0.50	0.10 - 0.40	0.18
Sodium Stearoyl Lactylate	0.30 - 3.0	0.50 - 2.50	1.00
Sodium Isostearoyl Lactylate	0.05 - 1.0	0.10 - 0.70	0.25
Tetrasodium EDTA	0.05 - 0.25	0.10 - 0.20	0.15
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	1.0 - 8.0	2.0 - 7.0	3.80
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Allantoin	0.50 - 2.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20

Example 5Preparation of Allantoin-Containing Skin Cream with Carboxypolymethylene Polymer

5

An allantoin-containing skin cream with carboxypolymethylene polymer is prepared according to Table 7. The preparation follows the method used in Example 3, except that the triethanolamine (Part D) is added last, after the combining of Parts A, B, and C, to avoid thickening of the emulsion. The triethanolamine is added to adjust the pH.

TABLE 7
ALLANTOIN-CONTAINING SKIN CREAM WITH
CARBOXYPOLYMETHYLENE POLYMER

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	60.0 - 80.0	73.55
Carboxypolymethylene Polymer	0.40 - 3.0	0.50 - 2.0	1.00
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.00
Cetyl Alcohol	1.0 - 8.0	2.0 - 7.0	3.00
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Allantoin	0.50 - 2.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20
<u>Part D</u>			
Triethanolamine (99%)	0.05 - 3.0	0.20 - 2.0	0.80

Example 6Preparation of Allantoin-Containing Skin Cream with Polyethylene Glycol Ethers of Cetearyl Alcohol

5

An allantoin-containing skin cream with polyethylene glycol ethers of cetearyl alcohol is prepared according to Table 8. The preparation follows the method used in Example 3. The citric acid is added to adjust the pH.

TABLE 8
ALLANTOIN-CONTAINING SKIN CREAM WITH
POLYETHYLENE GLYCOL ETHERS OF CETEARYL ALCOHOL

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	55.0 - 75.0	66.33
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
Tetrasodium EDTA	0.05 - 0.50	0.10 - 0.30	0.15
Ceteareth-25	0.50 - 4.0	2.00 - 3.50	2.60
Citric Acid	0.04 - 0.40	0.10 - 0.30	0.12
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	3.0 - 10.0	3.5 - 7.5	4.30
Stearyl Alcohol	1.0 - 5.0	2.0 - 4.0	3.50
Ceteareth-6	0.50 - 4.0	1.0 - 3.0	1.80
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Diazolidinyl Urea	0.05 - 0.50	0.10 - 0.30	0.15
Allantoin	0.50 - 2.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.30	0.20

Example 7Preparation of Allantoin-Containing Skin Cream with Polyethylene Glycol Ester of Stearic Acid
and Glyceryl Stearate

5

An allantoin-containing skin cream with a polyethylene glycol ester of stearic acid and glyceryl stearate is prepared according to Table 9. The preparation follows the method used in Example 3. The citric acid is added to adjust the pH.

TABLE 9
ALLANTOIN-CONTAINING SKIN CREAM WITH
POLYETHYLENE GLYCOL ESTER OF STEARIC ACID
AND GLYCERYL STEARATE

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	55.0 - 80.0	67.86
Propylene Glycol	2.0 - 9.0	4.3 - 7.0	5.70
Tetrasodium EDTA	0.05 - 0.50	0.10 - 0.30	0.15
Citric Acid	0.04 - 0.40	0.10 - 0.30	0.14
PEG-100 Stearate	1.0 - 5.0	1.5 - 3.0	2.60
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	2.0 - 12.0	10.60
Cetyl Alcohol	3.0 - 10.0	2.5 - 7.5	3.0
Stearyl Alcohol	1.0 - 4.0	1.0 - 3.5	2.50
Glyceryl Stearate	1.0 - 5.0	2.0 - 4.0	2.50
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Diazolidinyl Urea	0.05 - 0.50	0.10 - 0.30	0.20
Allantoin	0.50 - 2.00	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20

Example 8Preparation of Allantoin-Containing Skin Cream with Carboxypolymethylene Polymer and
Polyethylene Glycol Ester of Stearic Acid

5

An allantoin-containing skin cream with a carboxypolymethylene polymer and a polyethylene glycol ester of stearic acid is prepared according to Table 10. The preparation follows the method used in Example 5, with the triethanolamine (Part D) being added last. The triethanolamine is added to adjust the pH.

10

TABLE 10

ALLANTOIN-CONTAINING SKIN CREAM WITH
A CARBOXYPOLYMETHYLENE POLYMER AND A
POLYETHYLENE GLYCOL ESTER OF STEARIC ACID

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	60.0 - 85.0	69.95
Carboxypolymethylene Polymer	0.30 - 3.0	0.50 - 2.0	0.85
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
PEG-100 Stearate	0.25 - 2.5	0.50 - 2.0	1.50
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	1.0 - 8.0	2.0 - 7.0	4.20
Stearyl Alcohol	0.50 - 6.0	0.75 - 5.0	1.50
Cod Liver Oil	1.0 - 7.0	1.0 - 4.0	2.00
Butylated Hydroxytoluene	0.10 - 1.0	0.20 - 0.80	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Diazolidinyl Urea	0.05 - 0.25	0.10 - 0.20	0.15
Allantoin	0.50 - 2.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20
<u>Part D</u>			
Triethanolamine (99%)	0.05 - 3.0	0.20 - 2.0	0.80

Example 9Preparation of Allantoin-Containing Skin Cream with Galactoarabinan, Sodium Lauryl Sulfate,
and Beeswax

5

An allantoin-containing skin cream with galactoarabinan, sodium lauryl sulfate, and beeswax is prepared according to Table 11. The preparation follows the method used in Example 3. The citric acid is used to adjust the pH.

TABLE 11
ALLANTOIN-CONTAINING SKIN CREAM WITH
GALACTOARABINAN, SODIUM LAURYL SULFATE,
AND BEESWAX

<u>INGREDIENT</u>	<u>RANGE</u>	<u>PREFERRED</u>	<u>OPTIMUM</u>
<u>Part A</u>			
Water	50.0 - 90.0	60.0 - 80.0	61.65
Propylene Glycol	2.0 - 9.0	4.0 - 7.0	5.70
Sodium Lauryl Sulfate (30%)	0.50 - 5.0	1.0 - 3.0	1.90
Tetrasodium EDTA	0.05 - 0.30	0.10 - 0.20	0.15
Galactoarabinan	1.0 - 25.0	3.0 - 15.0	5.00
Citric Acid	0.05 - 0.25	0.10 - 0.20	0.15
<u>Part B</u>			
Lanolin Oil	5.0 - 15.0	8.0 - 12.0	10.60
Cetyl Alcohol	1.0 - 8.0	2.0 - 7.0	4.20
Stearyl Alcohol	0.50 - 6.0	1.0 - 4.0	2.00
Beeswax	0.50 - 5.0	1.0 - 3.0	1.90
Cod Liver Oil	0.50 - 15.0	1.0 - 10.0	2.00
Butylated Hydroxytoluene	0.10 - 3.0	0.25 - 2.5	0.50
<u>Part C</u>			
Methylparaben	0.10 - 0.50	0.15 - 0.40	0.30
Propylparaben	0.10 - 0.50	0.15 - 0.40	0.25
Allantoin	0.50 - 2.0	1.0 - 2.0	1.50
Fragrance	0.05 - 0.50	0.10 - 0.40	0.20

ADVANTAGES OF THE PRESENT INVENTION

5 The present invention provides an allantoin-containing composition that is an oil-
water emulsion using either an acidic anionic polymer and an anionic emulsifier or an acidic
anionic polymer and a nonionic emulsifier that is an ethoxylated ether or ethoxylated ester. The
composition can further include a carbohydrate polymer that can be polygalactose or
polyarabinose. If an ethoxylated ether or ethoxylated ester is used, the composition can further
10 comprise glyceryl stearate. The composition has improved thermal stability. The composition
according to the present invention is useful for treatment of a number of skin diseases and
conditions, including epidermolysis bullosa, pressure ulcers, diabetic ulcers, decubitus ulcers,
and milia. Compositions according to the present invention are also useful as skin protectants
even when these diseases are not present. Compositions according to the present invention are
15 well tolerated and can be used with other treatments.

 Although the present invention has been described in considerable detail, with
reference to certain preferred versions thereof, other versions and embodiments are possible.
Therefore, the scope of the invention is determined by the following claims.